

Jump to: [Page Content](#), [Section Navigation](#), [Site Navigation](#), [Site Search](#), [Account Information](#), or [Site Tools](#).

Note to users. If you're seeing this message, it means that your browser cannot find this page's style/presentation instructions -- or possibly that you are using a browser that does not support current Web standards. [Find out more](#) about why this message is appearing, and what you can do to make your experience of our site the best it can be.

Science/AAAS
Science
YAN JUN Alerts | Access Rights

AAAS.ORG Feedback Help
Librarians | My Account |

Advanced

AAAS

• News
NEWS

• Science Journals
SCIENCE JOURNALS

• Careers
CAREERS

• Communities
BLOGS & COMMUNITIES

• Multimedia
MULTIMEDIA

JOIN / SUBSCRIBE

Science The World's Journal of Original Scientific Research, Global News, and Commentary.

Science Home

Current Issue

Previous Issues

Science Express

Science Products

My Science

About the Journal

[Home](#) > [Science Magazine](#) > [17](#)

[July 1998](#) > Ronaghi et al. , pp. 363 - 365

Science 17 July 1998:
Vol. 281. no. 5375, pp. 363 - 365
DOI: 10.1126/science.281.5375.363

[Prev](#) | [Table of Contents](#) | [Next](#)

Tech.Sight

DNA SEQUENCING:

A Sequencing Method Based on Real-Time Pyrophosphate

Mostafa Ronaghi, Mathias Uhlén, and Pål Nyrén*

DNA sequencing is one of the most important technologies in bioscience today. Whole-genome approaches (1) and human expressed sequence tag (EST) sequencing (2) have started to exert profound influence on biology and medicine. The need for robust, high-throughput methods to replace the elegant Sanger method, described more than 20 years ago (3), has led to the development of several new principles, such as array methods based on sequencing by hybridization (4). New applications, such as population-based biodiversity projects and brute-force genotyping using single-nucleotide polymorphism, make such efforts even more urgent, in particular, for simple and robust methods for sequencing short "tags" (1 to 20 bases) such as ESTs or biallelic markers and methods suitable for routine diagnostic applications.

Sequencing-by-synthesis is based on the detection of nucleotide incorporation, using a primer-directed polymerase extension. The sequence can be deduced iteratively (5). During the last decade, many researchers, including the groups of Rosenthal (6), Gibbs (7), and Jones (8), described various protocols based on fluorescently labeled nucleotides. The level of incorporation of these fluorescent nucleotides is low, however, as shown by Metzker *et al.* (7), and therefore, the protocols only permit detection of a few bases.

Recently, Ronaghi *et al.* (9) showed that natural nucleotides can be used to obtain efficient incorporation during a sequencing-by-synthesis protocol. The detection was based on the pyrophosphate (PPi) released during the DNA polymerase reaction, the quantitative conversion of pyrophosphate to ATP by sulfurylase, and the subsequent production of visible light by firefly luciferase. However, this PPi-based sequencing method is not without drawbacks: The template must be washed thoroughly between nucleotide additions to remove unincorporated nucleotides. Also, templates not bound to a solid support are difficult to sequence, and the addition of new enzymes to each cycle of deoxynucleotide (dATP, dTTP, dGTP, and dCTP) is required.




Fig. 1. In the new DNA sequencing method, four nucleotides are added stepwise to the template hybridized to a primer. The PPi released in the DNA polymerase-catalyzed reaction is detected by the ATP sulfurylase and luciferase in a coupled reaction. The added nucleotides are continuously degraded by a nucleotide-degrading enzyme. After the first added nucleotide has been degraded, the next nucleotide can be added. As this procedure is repeated, longer stretches of the template sequence are deduced. dXTP, one of the four nucleotides.

Here, we address these problems by a modification in which the sequencing cycles can be performed without intermediate washing steps. This is achieved by the addition of a nucleotide-degrading enzyme to obtain a four-enzyme mixture. The principle of pyrosequencing is outlined in Fig. 1. The DNA fragment of interest (sequencing primer hybridized to a single-stranded DNA template) is incubated with DNA polymerase, ATP sulfurylase, firefly luciferase, and a nucleotide-degrading enzyme (such as apyrase). Repeated cycles of deoxynucleotide addition are performed. A deoxynucleotide will only be incorporated into the growing DNA strand if it is complementary to the base in the template strand. The synthesis of DNA is accompanied by release of PPi equal in molarity to that of the incorporated deoxynucleotide. Thereby, real-time signals are obtained by the enzymatic inorganic pyrophosphate detection assay (10). In this assay the released PPi is converted to ATP by ATP sulfurylase and the concentration of ATP is then sensed by the luciferase. The amount of light produced in the luciferase-catalyzed reaction can readily be estimated by a suitable light-sensitive device such as a luminometer or a CCD (charge-coupled device) camera. Unincorporated deoxynucleotides and the produced ATP are degraded between each cycle by the nucleotide-degrading enzyme. The nucleotide-degrading enzyme must possess the following properties: First, the enzyme must hydrolyze all deoxynucleotide triphosphate at approximately the same rate. This includes the α -thio-dATP, which is used instead of dATP to improve the background in sequencing reactions (9). Second, it should also hydrolyze ATP to prevent the accumulation of ATP between cycles. Third, the time for nucleotide degradation by the nucleotide-degrading enzyme must be slower than nucleotide incorporation by the polymerase. Obviously, these two enzymes compete for the same substrate, and it is important that the yield of primer-directed incorporation is as close to 100% as possible before the nucleotide-degrading enzyme can degrade the nucleotide to a concentration below the K_M for the polymerase. Finally, the rate of ATP synthesis by the sulfurylase should preferably be faster than the rate of ATP hydrolysis to obtain ATP concentrations and light production in proportion to the amount of PPi released.

To optimize the assay, several parameters were tested by using synthetic oligonucleotides as template (11). The assay solution consisted of the template-primer plus varying amounts of the four enzymes (12). The protocol consisted of simply adding a new nucleotide every other minute in an iterative manner and detecting the visible light produced. In Fig. 2 (Left), an example of the results is shown. Clear specific signals can be observed with low background. Note the higher signals obtained at cycles 21, 23, and 24 when two nucleotides were incorporated because of the presence of two identical adjacent bases in the template. Similar high-quality sequencing results were obtained for other oligonucleotide templates (13).




Fig. 2. Pyrosequencing was performed on a 35-base-long oligonucleotide template (**Left**) and a 130-base-long PCR product (**Right**). About 2 pmol of the template-primer was used in the assay. The reaction was started by the addition of the indicated deoxynucleotide, and the PPI released was detected by the described method. The DNA sequence after the primer is indicated.

For direct sequencing of polymerase chain reaction (PCR) products, single-stranded template was obtained by biotin capture on magnetic beads (14). In Fig. 2 (Right), the amount of light produced during approximately 40 subsequent cycles is shown. Sequence data covering 34 bases is obtained (15), and the signal-to-noise ratio remains relatively high even at 40 cycles. The low signals in the early cycles (2, 3, 6, and 7) show that the misincorporation of bases, which would produce PPI and consequently light, is not a major problem, despite the presence of "wrong" noncomplementary nucleotides. The background does increase during the later cycles, which is not surprising because relatively crude enzyme preparations were used in this work. Low amounts of contaminating enzymes such as exonucleases or kinases will give rise to nonsynchronized extensions on some templates, causing increased background and lower specific signal. Obviously, more work is needed to purify these enzymes from present contaminants. In addition, apyrase activity is decreased in later cycles, which is because of accumulation of intermediate products (such as deoxynucleoside diphosphate, or dNDP) and eventually undegraded dNTP. Removal of these nucleotides by enzymes such as nonspecific nucleoside diphosphatases (16) will increase the efficiency of apyrase in degradation of nucleoside triphosphates and thereby allow longer reads. However, it is reassuring that the nonoptimal enzyme mixture used here allows accurate determination of more than 20 bases for many different PCR products tested (13).

An inherent problem with the described method is the difficulty in determining the number of incorporated nucleotides in homopolymeric regions due to the nonlinear light response following incorporation of more than three or four identical nucleotides. This can be demonstrated by the relatively low signal at cycle 16 (Fig. 2, Right) when four T bases are incorporated. However, this nonlinear response can most likely be compensated for by software algorithms. In addition, for most tag-sequencing applications, such as brute-force EST-sequencing, biallelic marker analysis, and confirmatory sequencing, this problem is not a major concern, because the number of bases, if present, will be known.

With this method, parallel processing of large numbers of samples can easily be envisioned with the use of high-density microtiter plates and microinjector technology. An automated instrument has recently been developed based on the precise delivery of submicroliter volumes of the four nucleotides by "ink-jet" technology into a microtiter plate coupled with simultaneous detection of all samples by a single CCD unit (17). Together with a robot (17) performing single-strand template preparation (from double-stranded PCR products), ready for pyrosequencing, it would be possible to analyze thousands of samples daily with little manual intervention.

References and Notes

1. R. D. Fleischmann *et al.*, *Science* **269**, 496 (1995); C. M. Fraser *et al.*, *ibid.* **270**, 397 (1995); C. J. Bult *et al.*, *ibid.* **273**, 1058 (1996); J. Tomb *et al.*, *Nature* **388**, 539 (1997); H. W. Mewes *et al.*, *ibid.* **387**, 7 (1997).
2. V. E. Velculescu, L. Zhang, B. Vogelstein, K. W. Kinzler, *Science* **270**, 484 (1995).
3. F. Sanger, S. Nicklen, A. R. Coulson, *Proc. Natl. Acad. Sci. U.S.A.* **74**, 5463 (1977).
4. R. Drmanac, I. Labat, I. Brukner, R. Crkvenjakov, *Genomics* **4**, 114 (1989); H. Köster *et al.*, *Nature Biotechnol.* **14**, 1123 (1996).
5. E. D. Hyman, *Anal. Biochem.* **174**, 423 (1988).
6. A. Rosenthal, *International Patent Application Publication* 761107 (1989).
7. M. L. Metzker *et al.*, *Nucleic Acids Res.* **22**, 4259 (1994).
8. D. H. Jones, *Biotechniques* **22**, 938 (1997).
9. M. Ronaghi, S. Karamohamed, B. Pettersson, M. Uhlén, P. Nyrén, *Anal. Biochem.* **242**, 84 (1996).
10. P. Nyrén and A. Lundin, *ibid.* **151**, 504 (1985).
11. The oligonucleotides E3PN (5'-GCTGGAATTCGTCAGACTGGCCGTCGTTTTACAAC-3'), NUSPT (5'-GTAAAACGACGGCCAGT-3'), and JA80 (5'-GATGGAAACCAAAAATGATAGG-3') were synthesized by phosphoramidite chemistry (Interactiva).
12. The oligonucleotide E3PN and the PCR product generated from cloned HIV-V3 were used as templates for DNA sequencing. The oligonucleotides and single-stranded PCR product were hybridized to the primers NUSPT and JA80, respectively. The hybridized DNA fragments were incubated with *exo*⁻ Klenow or *exo*⁻

T7 DNA polymerase (Sequenase 2.0), respectively (Amersham). The sequencing procedure was carried out by stepwise elongation of the primer-strand upon sequential addition of the different deoxynucleoside triphosphates and simultaneous degradation of nucleotides by apyrase (nucleoside 5'-triphosphatase and nucleoside 5'-diphosphatase; EC 3.6.1.5) (Sigma). The sequencing reaction was performed at room temperature and was started by adding a specific amount of one of the deoxynucleotides. The PPi released due to nucleotide incorporation was detected as described (9).

13. M. Ronaghi and P. Nyrén, data not shown.
14. The biotinylated PCR products were immobilized onto streptavidin-coated super paramagnetic beads [Dynabeads M280-Streptavidin (Dyna)]. Elution of single-stranded DNA and hybridization of sequencing primers was carried out as described (9).
15. The sequencing data obtained from the pyrosequencing method was confirmed by semiautomated solid-phase Sanger sequencing (18).
16. H. D. Doremus and D. G. Belvins, *Plant Physiol.* **87**, 36 (1988).
17. The pyrosequencing instrument was based on a cassette containing the four separate nucleotides on an x-ray robotic arm (B. Ekström, M. Ronaghi, T. Nordström, P. Nyrén, M. Uhlén, unpublished data). The sample preparation robot was based on streptavidin-coated magnetic particles for PCR-capture and handling (A. Holmberg and M. Uhlén, unpublished data).
18. T. Hultman, S. Ståhl, E. Hornes, M. Uhlén, *Nucleic Acids Res.* **17**, 4937 (1989).
19. We thank K. Nourizad for very helpful technical assistance and A. Scott for critical review. Supported by grants from PyroSequencing AB and the Swedish Research Council for Engineering Science (TFR).

Mailbox: www.sciencemag.org/dmail.cgi?53491

The authors are in the Department of Biochemistry and Biotechnology, The Royal Institute of Technology, SE-10044 Stockholm, Sweden. E-mail: paaln@biochem.kth.se

**To whom correspondence should be addressed.*

THIS ARTICLE HAS BEEN CITED BY OTHER ARTICLES:

Pyrosequencing for Rapid Detection of Mycobacterium tuberculosis Resistance to Rifampin, Isoniazid, and Fluoroquinolones.

L. T. C. Bravo, M. J. Tuohy, C. Ang, R. V. Destura, M. Mendoza, G. W. Procop, S. M. Gordon, G. S. Hall, and N. K. Shrestha (2009)

J. Clin. Microbiol. **47**, 3985-3990

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Rapid identification of homologous recombinants and determination of gene copy number with reference/query pyrosequencing (RQPS).

Z. Liu, A. C. Obenauf, M. R. Speicher, and R. Kopan (2009)

Genome Res. **19**, 2081-2089

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

PerM: efficient mapping of short sequencing reads with periodic full sensitive spaced seeds.

Y. Chen, T. Souaiaia, and T. Chen (2009)

Bioinformatics **25**, 2514-2521

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Orphelia: predicting genes in metagenomic sequencing reads.

K. J. Hoff, T. Lingner, P. Meinicke, and M. Tech (2009)

Nucleic Acids Res. **37**, W101-W105

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Flow cytometry for enrichment and titration in massively parallel DNA sequencing.

J. Sandberg, P. L. Stahl, A. Ahmadian, M. K. Bjursell, and J. Lundeberg (2009)

Nucleic Acids Res. **37**, e63

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Whole-Genome-Based Phylogeny and Divergence of the Genus *Brucella*.

J. T. Foster, S. M. Beckstrom-Sternberg, T. Pearson, J. S. Beckstrom-Sternberg, P. S. G. Chain, F. F.

Roberto, J. Hnath, T. Brettin, and P. Keim (2009)

J. Bacteriol. **191**, 2864-2870

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Next-Generation Sequencing: From Basic Research to Diagnostics.

K. V. Voelkerding, S. A. Dames, and J. D. Durtschi (2009)

Clin. Chem. **55**, 641-658

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Real-Time DNA Sequencing from Single Polymerase Molecules.

J. Eid, A. Fehr, J. Gray, K. Luong, J. Lyle, G. Otto, P. Peluso, D. Rank, P. Baybayan, B. Bettman, *et al.* (2009)

Science **323**, 133-138

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

The perils of plenty: what are we going to do with all these genes?.

A. Rodrigo, F. Bertels, J. Heled, R. Noder, H. Shearman, and P. Tsai (2008)

Phil Trans R Soc B **363**, 3893-3902

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Pyrosequencing analysis of the Oral Microflora of healthy adults.

B.J.F. Keijser, E. Zaura, S.M. Huse, J.M.B.M. van der Vossen, F.H.J. Schuren, R.C. Montijn, J.M. ten Cate, and W. Crielaard (2008)

Journal of Dental Research **87**, 1016-1020

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Nested Patch PCR enables highly multiplexed mutation discovery in candidate genes.

K. E. Varley and R. D. Mitra (2008)

Genome Res. **18**, 1844-1850

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Cerebellar {alpha}-synuclein levels are decreased in Parkinson's disease and do not correlate with SNCA polymorphisms associated with disease in a Swedish material.

M. Westerlund, A. C. Belin, A. Anvret, A. Hakansson, H. Nissbrandt, C. Lind, O. Sydow, L. Olson, and D. Galter (2008)

FASEB J **22**, 3509-3514

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Four-color DNA sequencing with 3'-O-modified nucleotide reversible terminators and chemically cleavable fluorescent dideoxynucleotides.

J. Guo, N. Xu, Z. Li, S. Zhang, J. Wu, D. H. Kim, M. Sano Marma, Q. Meng, H. Cao, X. Li, *et al.* (2008)

PNAS **105**, 9145-9150

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Typing of SHV Extended-Spectrum {beta}-Lactamases by Pyrosequencing in *Klebsiella pneumoniae* Strains with Chromosomal SHV {beta}-Lactamase.

M. Haanpera, S. D. Forssten, P. Huovinen, and J. Jalava (2008)

Antimicrob. Agents Chemother. **52**, 2632-2635

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

The new paradigm of flow cell sequencing.

R. A. Holt and S. J.M. Jones (2008)

Genome Res. **18**, 839-846

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Frequency and Effect of the Bovine Acyl-CoA:Diacylglycerol Acyltransferase 1 (DGAT1) K232A Polymorphism in Swedish Dairy Cattle.

J. Naslund, W. F. Fikse, G. R. Pielberg, and A. Lunden (2008)

J Dairy Sci **91**, 2127-2134

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Rapid comparative genomic analysis for clinical microbiology: The *Francisella tularensis* paradigm.

B. La Scola, K. Elkarkouri, W. Li, T. Wahab, G. Fournous, J.-M. Rolain, S. Biswas, M. Drancourt, C. Robert, S. Audic, *et al.* (2008)

Genome Res. **18**, 742-750

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Quantitative Promoter Hypermethylation Analysis of Cancer-Related Genes in Salivary Gland Carcinomas: Comparison with Methylation-Specific PCR Technique and Clinical Significance.

E.-S. Lee, J.-P. Issa, D. B. Roberts, M. D. Williams, R. S. Weber, M. S. Kies, and A. K. El-Naggar (2008)

Clin. Cancer Res. **14**, 2664-2672

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

The HIV-1 protease substitution K55R: a protease-inhibitor-associated substitution involved in restoring viral replication.

E. S. Margerison, M. Maguire, D. Pillay, P. Cane, and R. C. Elston (2008)

J. Antimicrob. Chemother. **61**, 786-791

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Combined Real-Time PCR and Pyrosequencing Strategy for Objective, Sensitive, Specific, and High-Throughput Identification of Reduced Susceptibility to Penicillins in *Neisseria meningitidis*.

S. Thulin, P. Olcen, H. Fredlund, and M. Unemo (2008)

Antimicrob. Agents Chemother. **52**, 753-756

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Short read fragment assembly of bacterial genomes.

M. J. Chaisson and P. A. Pevzner (2008)

Genome Res. **18**, 324-330

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Multiple Hybridization-extension Sequencing (MHES) on Microarray.

Z. Pan, Y. Li, P. Xiao, and Z. Lu (2007)

J. Biochem. **142**, 605-611

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

3'-O-modified nucleotides as reversible terminators for pyrosequencing.

J. Wu, S. Zhang, Q. Meng, H. Cao, Z. Li, X. Li, S. Shi, D. H. Kim, L. Bi, N. J. Turro, *et al.* (2007)

PNAS **104**, 16462-16467

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Termination of DNA synthesis by N6-alkylated, not 3'-O-alkylated, photocleavable 2'-deoxyadenosine triphosphates.

W. Wu, B. P. Stupi, V. A. Litosh, D. Mansouri, D. Farley, S. Morris, S. Metzker, and M. L. Metzker (2007)

Nucleic Acids Res. **35**, 6339-6349

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Plasma Vascular Endothelial Growth Factor-A (VEGF-A) and VEGF-A Gene Polymorphism are Associated with Hydrocele Development in Lymphatic Filariasis.

A. Y. Debrah, S. Mand, M. R. Toliat, Y. Marfo-Debrekyei, L. Batsa, P. Nurnberg, B. Lawson, O. Adjei, A. Hoerauf, and K. Pfarr (2007)

Am J Trop Med Hyg **77**, 601-608

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Gene-environment interactions in parkinsonism and Parkinson's disease: the Geoparkinson study.

F D Dick, G De Palma, A Ahmadi, A Osborne, N W Scott, G J Prescott, J Bennett, S Semple, S Dick, P Mozzoni, *et al.* (2007)

Occup. Environ. Med. **64**, 673-680

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Rapid screening of clarithromycin resistance in *Helicobacter pylori* by pyrosequencing.

K.-A. Moder, F. Layer, W. König, and B. König (2007)

J. Med. Microbiol. **56**, 1370-1376

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Article Views

- [Summary](#)
- Full Text (HTML)

Article Tools

- [Save to My Folders](#)
- [Download Citation](#)
- [Alert Me When Article is Cited](#)
- [Post to CiteULike](#)
- [E-mail This Page](#)
- [Submit an E-Letter](#)
- [Commercial Reprints and Permissions](#)
- [View PubMed Citation](#)

Related Content

Similar Articles In:

- [Science Magazine](#)
- [Web of Science](#)
- [PubMed](#)

Search Google Scholar for:

- [Articles by Ronaghi, M.](#)
- [Articles by Nyberg, a. P.](#)

Find Citing Articles in:

- [Web of Science \(466\)](#)
- [HighWire Press](#)
- [CrossRef](#)

Rapid Identification of Promoter Hypermethylation in Hepatocellular Carcinoma by Pyrosequencing of Etiologically Homogeneous Sample Pools.

E. Dejeux, V. Audard, C. Cavard, I. G. Gut, B. Terris, and J. Tost (2007)

J. Mol. Diagn. **9**, 510-520

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Multigene amplification and massively parallel sequencing for cancer mutation discovery.

F. Dahl, J. Stenberg, S. Fredriksson, K. Welch, M. Zhang, M. Nilsson, D. Bicknell, W. F. Bodmer, R. W. Davis, and H. Ji (2007)

PNAS **104**, 9387-9392

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

More on Contamination: The Use of Asymmetric Molecular Behavior to Identify Authentic Ancient Human DNA.

H. Malmstrom, E. M. Svensson, M. T. P. Gilbert, E. Willerslev, A. Götherström, and G. Holmlund (2007)

Mol. Biol. Evol. **24**, 998-1004

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Identification of Alpha-Hemolytic Streptococci by Pyrosequencing the 16S rRNA Gene and by Use of VITEK 2.

M. Haanpera, J. Jalava, P. Huovinen, O. Meurman, and K. Rantakokko-Jalava (2007)

J. Clin. Microbiol. **45**, 762-770

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Four-color DNA sequencing by synthesis using cleavable fluorescent nucleotide reversible terminators.

J. Ju, D. H. Kim, L. Bi, Q. Meng, X. Bai, Z. Li, X. Li, M. S. Marma, S. Shi, J. Wu, *et al.* (2006)

PNAS **103**, 19635-19640

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

A Sanger/pyrosequencing hybrid approach for the generation of high-quality draft assemblies of marine microbial genomes.

S. M. D. Goldberg, J. Johnson, D. Busam, T. Feldblyum, S. Ferreira, R. Friedman, A. Halpern, H. Khouri, S. A. Kravitz, F. M. Lauro, *et al.* (2006)

PNAS **103**, 11240-11245

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Rapid Detection of Rifampin Resistance in *Mycobacterium tuberculosis* by Pyrosequencing Technology..

P. Jureen, L. Engstrand, S. Eriksson, A. Alderborn, M. Krabbe, and S. E. Hoffner (2006)

ADVERTISEMENT



ADVERTISEMENT

Introducing:

The new journal
from AAAS
& Science.

At the expanding
intersection of basic science
and clinical medicine.

Click Here for
more information
and to subscribe

- [Google Scholar](#)
- [Scopus](#)

J. Clin. Microbiol. **44**, 1925-1929
 | [Abstract »](#) | [Full Text »](#) | [PDF »](#)

My Science

- [My Folders](#)
- [My Alerts](#)
- [My Saved Searches](#)
- [Sign Out](#)

Quantitative Analysis of SRNPN Gene Methylation by Pyrosequencing as a Diagnostic Test for Prader-Willi Syndrome and Angelman Syndrome.

H. E. White, V. J. Durston, J. F. Harvey, and N. C.P. Cross (2006)
Clin. Chem. **52**, 1005-1013
 | [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Streptococcus pneumoniae Isolates Resistant to Telithromycin..

M. Rantala, M. Haanpera-Heikkinen, M. Lindgren, H. Seppala, P. Huovinen, J. Jalava, and the Finnish Study Group for Antimicrobial Resistan (2006)
Antimicrob. Agents Chemother. **50**, 1855-1858
 | [Abstract »](#) | [Full Text »](#) | [PDF »](#)

DNA sequence variation in the promoter region of the VEGF gene impacts VEGF gene expression and maximal oxygen consumption.

S. J. Prior, J. M. Hagberg, C. M. Paton, L. W. Douglass, M. D. Brown, J. C. McLenithan, and S. M. Roth (2006)
Am J Physiol Heart Circ Physiol **290**, H1848-H1855
 | [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Emerging technologies in DNA sequencing.

M. L. Metzker (2005)
Genome Res. **15**, 1767-1776
 | [Abstract »](#) | [Full Text »](#) | [PDF »](#)

An analysis of the feasibility of short read sequencing.

N. Whiteford, N. Haslam, G. Weber, A. Prugel-Bennett, J. W. Essex, P. L. Roach, M. Bradley, and C. Neylon (2005)
Nucleic Acids Res. **33**, e171
 | [Abstract »](#) | [Full Text »](#) | [PDF »](#)

New Quinolone Resistance Phenomenon in Salmonella enterica: Nalidixic Acid-Susceptible Isolates with Reduced Fluoroquinolone Susceptibility.

A. J. Hakanen, M. Lindgren, P. Huovinen, J. Jalava, A. Siitonen, and P. Kotilainen (2005)
J. Clin. Microbiol. **43**, 5775-5778
 | [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Prevalence and Molecular Genetics of Macrolide Resistance among Streptococcus pneumoniae Isolates Collected in Finland in 2002.

M. Rantala, S. Huikko, P. Huovinen, J. Jalava, and the Finnish Study Group for Antimicrobial Resistan (2005)
Antimicrob. Agents Chemother. **49**, 4180-4184



[To Advertise](#) | [Find Products](#)

ADVERTISEMENT

Featured Jobs

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Estimating RNA editing efficiency of five editing sites in the serotonin 2C receptor by pyrosequencing.

K. IWAMOTO, M. BUNDO, and T. KATO (2005)

RNA **11**, 1596-1603

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Multiplex SNP typing by bioluminometric assay coupled with terminator incorporation (BATI).

G.-H. Zhou, M. Gotou, T. Kajiyama, and H. Kambara (2005)

Nucleic Acids Res. **33**, e133

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Discovery of SNPs in Soybean Genotypes Frequently Used as the Parents of Mapping Populations in the United States and Korea.

K. Van, E.-Y. Hwang, M. Y. Kim, H. J. Park, S.-H. Lee, and P. B. Cregan (2005)

J. Hered. **96**, 529-535

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Two-Step Genetic Screening of Thrombophilia by Pyrosequencing.

A. Verri, F. Focher, G. Tettamanti, and V. Grazioli (2005)

Clin. Chem. **51**, 1282-1284

| [Full Text »](#) | [PDF »](#)

Prehistoric contacts over the Straits of Gibraltar indicated by genetic analysis of Iberian Bronze Age cattle.

C. Anderung, A. Bouwman, P. Persson, J. M. Carretero, A. I. Ortega, R. Elburg, C. Smith, J. L. Arsuaga, H. Ellegren, and A. Gotherstrom (2005)

PNAS **102**, 8431-8435

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Four-color DNA sequencing by synthesis on a chip using photocleavable fluorescent nucleotides.

T. S. Seo, X. Bai, D. H. Kim, Q. Meng, S. Shi, H. Ruparel, Z. Li, N. J. Turro, and J. Ju (2005)

PNAS **102**, 5926-5931

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Competitive enzymatic reaction to control allele-specific extensions.

E. Hultin, M. Kaller, A. Ahmadian, and J. Lundeberg (2005)

Nucleic Acids Res. **33**, e48

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Determination of CYP2D6 Gene Copy Number by Pyrosequencing.

E. Soderback, A.-L. Zackrisson, B. Lindblom, and A. Alderborn (2005)

Clin. Chem. **51**, 522-531

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Detection of *Aspergillus fumigatus* and a Mutation That Confers Reduced Susceptibility to Itraconazole and Posaconazole by Real-Time PCR and Pyrosequencing.

J. P. Trama, E. Mordechai, and M. E. Adelson (2005)

J. Clin. Microbiol. **43**, 906-908

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Detection and Quantification of Macrolide Resistance Mutations at Positions 2058 and 2059 of the 23S rRNA Gene by Pyrosequencing.

M. Haanpera, P. Huovinen, and J. Jalava (2005)

Antimicrob. Agents Chemother. **49**, 457-460

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

A multi-enzyme model for pyrosequencing.

A. Agah, M. Aghajan, F. Mashayekhi, S. Amini, R. W. Davis, J. D. Plummer, M. Ronaghi, and P. B. Griffin (2004)

Nucleic Acids Res. **32**, e166

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Ceruloplasmin gene variations and substantia nigra hyperechogenicity in Parkinson disease.

H. Hochstrasser, P. Bauer, U. Walter, S. Behnke, J. Spiegel, I. Csoti, B. Zeiler, A. Bornemann, J. Pahnke, G. Becker, *et al.* (2004)

Neurology **63**, 1912-1917

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Analytical Validation of the Tag-It High-Throughput Microsphere-Based Universal Array Genotyping Platform: Application to the Multiplex Detection of a Panel of Thrombophilia-Associated Single-Nucleotide Polymorphisms.

S. Bortolin, M. Black, H. Modi, I. Boszko, D. Kobler, D. Fieldhouse, E. Lopes, J.-M. Lacroix, R. Grimwood, P. Wells, *et al.* (2004)

Clin. Chem. **50**, 2028-2036

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Pyrosequencing for Detection of Lamivudine-Resistant Hepatitis B Virus.

A. Lindstrom, J. Odeberg, and J. Albert (2004)

J. Clin. Microbiol. **42**, 4788-4795

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Characterization of synthetic DNA bar codes in *Saccharomyces cerevisiae* gene-deletion strains.

R. G. Eason, N. Pourmand, W. Tongprasit, Z. S. Herman, K. Anthony, O. Jejelowo, R. W. Davis, and V. Stolc (2004)

PNAS **101**, 11046-11051

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Molecular Typing of *Neisseria gonorrhoeae* Isolates by Pyrosequencing of Highly Polymorphic Segments of the

porB Gene.

M. Unemo, P. Olcen, J. Jonasson, and H. Fredlund (2004)

J. Clin. Microbiol. **42**, 2926-2934

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Microfluidic device reads up to four consecutive base pairs in DNA sequencing-by-synthesis.

E. P. Kartalov and S. R. Quake (2004)

Nucleic Acids Res. **32**, 2873-2879

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Sequence variation and conservation in virulence-related genes of *Bordetella pertussis* isolates from the UK.

E. R. Packard, R. Parton, J. G. Coote, and N. K. Fry (2004)

J. Med. Microbiol. **53**, 355-365

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Photocleavable fluorescent nucleotides for DNA sequencing on a chip constructed by site-specific coupling chemistry.

T. S. Seo, X. Bai, H. Ruparel, Z. Li, N. J. Turro, and J. Ju (2004)

PNAS **101**, 5488-5493

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Discrimination between Scrapie and Bovine Spongiform Encephalopathy in Sheep by Molecular Size, Immunoreactivity, and Glycoprofile of Prion Protein.

C. M. A. Thuring, J. H. F. Erkens, J. G. Jacobs, A. Bossers, L. J. M. Van Keulen, G. J. Garssen, F. G. Van Zijderveld, S. J. Ryder, M. H. Groschup, T. Sweeney, *et al.* (2004)

J. Clin. Microbiol. **42**, 972-980

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Digital Detection of Genetic Mutations Using SPC-Sequencing.

H. Ruparel, M. E. Ulz, S. Kim, and J. Ju (2004)

Genome Res. **14**, 296-300

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Rapid Detection and Estimation by Pyrosequencing of 23S rRNA Genes with a Single Nucleotide Polymorphism Conferring Linezolid Resistance in Enterococci.

A. Sinclair, C. Arnold, and N. Woodford (2003)

Antimicrob. Agents Chemother. **47**, 3620-3622

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Comparison of Serologic and Genetic porB-Based Typing of *Neisseria gonorrhoeae*: Consequences for Future Characterization.

M. Unemo, P. Olcen, J. Albert, and H. Fredlund (2003)

J. Clin. Microbiol. **41**, 4141-4147

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Single base extension (SBE) with proofreading polymerases and phosphorothioate primers: improved fidelity in single-substrate assays.

D. Di Giusto and G. C. King (2003)

Nucleic Acids Res. **31**, e7

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

A photocleavable fluorescent nucleotide for DNA sequencing and analysis.

Z. Li, X. Bai, H. Ruparel, S. Kim, N. J. Turro, and J. Ju (2003)

PNAS **100**, 414-419

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

NotI passorting to identify species composition of complex microbial systems.

V. Zabarovska, A. S. Kutsenko, L. Petrenko, G. Kilosanidze, O. Ljungqvist, E. Norin, T. Midtvedt, G. Winberg, R. Mollby, V. I. Kashuba, *et al.* (2003)

Nucleic Acids Res. **31**, e5

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

A Nonsense Mutation in the FMO3 Gene Underlies Fishy Off-Flavor in Cow's Milk.

A. Lunden, S. Marklund, V. Gustafsson, and L. Andersson (2002)

Genome Res. **12**, 1885-1888

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

hairy: A Quantitative Trait Locus for Drosophila Sensory Bristle Number.

C. Robin, R. F. Lyman, A. D. Long, C. H. Langley, and T. F. C. Mackay (2002)

Genetics **162**, 155-164

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Bioluminescent Method for Detecting Telomerase Activity.

S.-Q. Xu, M. He, H.-P. Yu, X.-Y. Wang, X.-L. Tan, B. Lu, X. Sun, Y.-K. Zhou, Q.-F. Yao, Y.-J. Xu, *et al.* (2002)

Clin. Chem. **48**, 1016-1020

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Infusion of CD4+ Donor Lymphocytes Induces the Expansion of CD8+ Donor T Cells with Cytolytic Activity Directed against Recipient Hematopoietic Cells.

E. Zorn, K. S. Wang, E. P. Hochberg, C. Canning, E. P. Alyea, R. J. Soiffer, and J. Ritz (2002)

Clin. Cancer Res. **8**, 2052-2060

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Multiplex Pyrosequencing.

N. Pourmand, E. Elahi, R. W. Davis, and M. Ronaghi (2002)

Nucleic Acids Res. **30**, e31

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

A new approach to genome mapping and sequencing: slalom libraries.

V. I. Zabarovska, R. Z. Gizatullin, A. N. Al-Amin, R. Podowski, A. I. Protopopov, S. Lofdahl, C.

Wahlestedt, G. Winberg, V. I. Kashuba, I. Ernberg, *et al.* (2002)

Nucleic Acids Res. **30**, e6

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Unexpectedly High Allelic Diversity at the KIT Locus Causing Dominant White Color in the Domestic Pig.

G. Pielberg, C. Olsson, A.-C. Syvanen, and L. Andersson (2002)

Genetics **160**, 305-311

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Genotyping by apyrase-mediated allele-specific extension.

A. Ahmadian, B. Gharizadeh, D. O'Meara, J. Odeberg, and J. Lundeberg (2001)

Nucleic Acids Res. **29**, e121

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

L-RCA (ligation-rolling circle amplification): a general method for genotyping of single nucleotide polymorphisms (SNPs).

X. Qi, S. Bakht, K. M. Devos, M. D. Gale, and A. Osbourn (2001)

Nucleic Acids Res. **29**, e116

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Pyrosequencing as a Method for Grouping of *Listeria monocytogenes* Strains on the Basis of Single-Nucleotide Polymorphisms in the *inlB* Gene.

H. Unnerstad, H. Ericsson, A. Alderborn, W. Tham, M.-L. Danielsson-Tham, and J. G. Mattsson (2001)

Appl. Envir. Microbiol. **67**, 5339-5342

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Quantitative detection of single nucleotide polymorphisms for a pooled sample by a bioluminometric assay coupled with modified primer extension reactions (BAMPER).

G.-h. Zhou, M. Kamahori, K. Okano, G. Chuan, K. Harada, and H. Kambara (2001)

Nucleic Acids Res. **29**, e93

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Monitoring Resistance to Human Immunodeficiency Virus Type 1 Protease Inhibitors by Pyrosequencing.

D. O'Meara, K. Wilbe, T. Leitner, B. Hejdeman, J. Albert, and J. Lundeberg (2001)

J. Clin. Microbiol. **39**, 464-473

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Pyrosequencing Sheds Light on DNA Sequencing.

M. Ronaghi (2001)

Genome Res. **11**, 3-11

| [Abstract »](#) | [Full Text »](#)

Automation for Genomics, Part Two: Sequencers, Microarrays, and Future Trends.

D. Meldrum (2000)

Genome Res. **10**, 1288-1303

| [Abstract »](#) | [Full Text »](#)

Determination of Single-Nucleotide Polymorphisms by Real-time Pyrophosphate DNA Sequencing.

A. Alderborn, A. Kristofferson, and U. Hammerling (2000)

Genome Res. **10**, 1249-1258

| [Abstract »](#) | [Full Text »](#)

A Mutation in PRKAG3 Associated with Excess Glycogen Content in Pig Skeletal Muscle.

D. Milan, J. Jeon, C. Looft, V. Amarger, A. Robic, M. Thelander, C. Rogel-Gaillard, S. Paul, N.

Iannuccelli, L. Rask, *et al.* (2000)

Science **288**, 1248-1251

| [Abstract »](#) | [Full Text »](#)

Production by quantitative photolithographic synthesis of individually quality checked DNA microarrays.

M. Beier and J. D. Hoheisel (2000)

Nucleic Acids Res. **28**, e11

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Polymorphism in the Pertussis Toxin Promoter Region Affecting the DNA-Based Diagnosis of Bordetella Infection.

M. Nygren, E. Reizenstein, M. Ronaghi, and J. Lundeberg (2000)

J. Clin. Microbiol. **38**, 55-60

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Molecular grouping of *Listeria monocytogenes* based on the sequence of the *inlB* gene.

H. ERICSSON, H. UNNERSTAD, J. G. MATTSSON, M.-L. DANIELSSON-THAM, and W. THAM (2000)

J. Med. Microbiol. **49**, 73-80

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)

Gene discovery in the wood-forming tissues of poplar: Analysis of 5,692 expressed sequence tags.

F. Sterky, S. Regan, J. Karlsson, M. Hertzberg, A. Rohde, A. Holmberg, B. Amini, R. Bhalerao, M.

Larsson, R. Villarroel, *et al.* (1998)

PNAS **95**, 13330-13335

| [Abstract »](#) | [Full Text »](#) | [PDF »](#)



[News](#) | [Science Journals](#) | [Careers](#) | [Blogs and Communities](#) | [Multimedia](#) | [Collections](#) | [Help](#) | [Site Map](#) | [RSS](#)

[Subscribe](#) | [Feedback](#) | [Privacy / Legal](#) | [About Us](#) | [Advertise With Us](#) | [Contact Us](#)

© 1998 American Association for the Advancement of Science. All Rights Reserved.

AAAS is a partner of [HINARI](#), [AGORA](#), [PatientInform](#), [CrossRef](#), and [COUNTER](#).

You have reached the bottom of the page. [Back to top](#)